Improving Human Nutrition Through Agriculture: The Role of International Agricultural Research

October 5-7, 1999

Quality Protein Maize Story

S.K. Vasal

CIMMYT

A Workshop hosted by the International Rice Research Institute, Los Banos, Philippines organized by the
International Food Policy Research Institute

This paper is intended to present a critique of various aspects of protein related nutritional improvements in maize. Earlier efforts and difficulties in developing germplasm products that failed to meet expectations and lacked agronomic performance and acceptance at the farmer's level will be discussed. Sequence of events will be presented that led to choosing an option and a strategy that turned out to be successful and has proved viable in developing a speciality corn type which is referred to as Quality Protein Maize (QPM). It has some characteristics different from original high lysine soft opaque-2 maize cultivars. It is distinct as it is no more opaque and is indistinguishable from normal endosperm maize types. Also, unlike soft opaques, QPM has been developed using a combined interaction of opaque-2 gene and the genetic modifiers while still maintaining superior quality protein.

Maize is world's one of the three most popular cereal crops. It is grown worldwide on approximately 130 million ha annually with a production of 574 million metric tons (Ito 1998). It occupies an important position in world economy and trade as a food, feed and an industrial grain crop. Several million people in the developing world consume maize as an important staple food and derive their protein and calories requirements from it. Maize is thus a potential source of protein for humans and animals. It holds a great promise for increasing production and also as a future protein crop. Genetic manipulation for improved nutritional value particularly protein content and quality is a noble goal. Realizing achievements through normal breeding efforts are difficult, complex, frustrating and require long-term investments, sustained research efforts, patience and continuing administrative, financial and scientific support. Penalty in yield and in other economic traits is not accepted unless the product has a value added properties.

Poor nutritional value of maize grain is well known and the need to improve it has been recognized for a long time (Osborne and Mendel, 1914). Most of the protein in a mature maize kernel is contained in the endosperm and the germ. Both parts differ substantially in the content and the quality of the protein. The endosperm protein is low in quantity as well as quality. In contrast the germ protein is sufficiently superior in quantity and quality. Since endosperm constitutes bulk of the grain, it is estimated it may contribute as much as 80% of the total kernel protein. Any major improvements for such trait(s) will thus have to be done in this part. Genetic variation for most traits in maize is incredibly high and amenable to enhancements. Time factor is major limitation requiring improvements extended over an unusually long period of time. Breeders and nutritionists alike agree that enhanced levels of both protein content and quality will greatly result in improved nutritional value of this grain. Attempts to improve protein content began towards the later part of the 19th century. The quality, however, became important and a concern at a much later date. Numerous researchers have attempted to improve protein content with good results. For more information the readers may consult relevant papers published on this subject. In this paper the focus will be only on improving protein quality presenting a critique and a detailed discussion of various aspects.

As is well known, serious efforts to improve the nutritional quality of maize endosperm protein began in the mid-60s. Prior to this period, however, the poor nutritional quality of maize was known and documented. Also efforts were rather limited

to only screening elite maize germplasm and accessions to identify genotypes superior for this trait. In the absence of specific gene(s), the alternate improvement strategy involving recurrent selection could not be used easily. It required step by step accumulation of favorable alleles for an extended period before a noticeable change in nutritional value could be observed. The lack of a simple genetic system also prevented use of a straightforward backcross program. Thus protein quality remained more of a concern but with no immediate solutions in sight. No action-oriented strategies could be deployed to resolve this issue. It may be important to study in depth the inherent causes responsible for poor protein quality of maize endosperm protein. The bulk of maize endosperm protein is comprised of zein fraction that is completely devoid of two important essential amino acids lysine and tryptophan. The amounts may vary but this fraction may constitute as much as 60% of the total protein. The high level of this particular fraction is the sole cause of poor protein quality. It is thus understandable that this fraction will have to be reduced substantially if improvements in nutritional characteristics are to be realized. In normal maize, the proportion of various fractions on the average is albumins (3%), globulins (3%), Zein (60%), and glutelin (34%). Further it may be pointed out that all fractions other than zein are balanced and quite rich in lysine and tryptophan. Suppression of lysine-deficient zein fraction without altering contribution of other fractions could thus be seen as a feasible approach to bring about improvements in the nutritional value of maize grain. A reduction in zein fraction causes proportional elevation of other fractions rich in lysine thus resulting in elevation of these two amino acids in protein but not on absolute basis of per unit of endosperm in the grain.

Beginning of genetic manipulation of protein quality

The discovery of the biochemical effects by the Purdue university researchers of mutant alleles opaque-2 (o2) and floury-2 (fl2) opened exciting opportunities thus paving the way for improving protein quality of maize endosperm protein (Mertz et al. 1964, Nelson et al. 1965). This much needed discovery aroused considerable interest, enthusiasm, and hopes worldwide. These mutants alter amino acid profile and composition of maize endosperm protein and result in two-fold tryptophan increase in the levels of lysine and tryptophan compared to what is encountered in normal genotypes. The mutants derive their name from soft floury opaque endosperm. The lysine in maize is the first limiting amino acid and tryptophan as the second. In addition other amino acids such as histidine, arginine, aspartic acid, and glycine also show an increase. A decline is observed in other amino acids such as glutamic acid, alanine and leucine. A decrease in leucine is considered desirable as it makes leucine-isoleucine ratio more balanced which in turn helps to liberate more tryptophan for more niacin biosynthesis and thus helping to combat pellagra. A sulfur containing amino acid methionine registers increase in only floury -2 mutant.

Besides biochemical traits, the mutants adversely affect agronomic performance including yield and kernel characteristics. The lower yield results from reduced dry matter accumulation. The kernel appearance is altered to soft chalky phenotype which is unattractive and not liked by maize growers in the developing countries. Physiological drying is also affected and a higher vulnerability is observed to ear rot causing organisms and stored grain insects pests. Variable genetic background effects are also noticed for such traits as pericarp thickness, larger germ size, reduced cob weight, seed color intensity, and kernel weight and density. Occurrence of pleiotropic effects facilitate

selection process in the desired direction. Genetic background effects are indeed important to harness beneficial effects of the mutant alleles and to remedy some of the drawbacks. Some of the effects may be environmentally specific as is the case with low germination in cold conditions. For a more detailed discussion on effects and causes the readers may refer to published literature on this subject (Harpstead, 1969; Bjarnason and Vasal, 1992).

A continuing search for identifying new and better mutants has been underway during the past several decades. In addition to opaque-2 and floury-2, other mutants have been discovered such as opaque-7(o7), opaque-6(o6), floury-3(fl3), mucronate and defective endosperm. Unfortunately up to this point no other mutant has been found which offers any additional advantage(s) over the opaque-2 gene in practical maize breeding programs. Several of the reported mutants have been experimentally tried at CIMMYT but the main focus on developing QPM germplasm has been on intensive use of the opaque-2 gene.

Early efforts and experience

New genes and gene combinations affecting nutritional improvements have been of interest and value to maize breeders worldwide. Unusual enthusiasm and excitement draw researcher's attention for possible use and exploitation of these traits. Same is true for the high lysine mutants in maize. Beginning mid-60s, improving protein quality became possible in developed developing countries. Elite germplasm and elite inbreds making good hybrid combinations were converted as rapidly as possible through standard backcross approach. In the initial stages both o-2 and fl-2 genes were used singly or in combination with each other. Later as some undesirable effects of fl-2 gene were discovered, its use slowed down and discontinued. It is fair to say that almost for a decade, the major emphasis in most breeding programs was on conversion programs to obtain mutant versions of normal genotypes. Many important commercial OPVs and hybrids were experimentally tested and exploited in farmer's fields. By the mid-70s enough experience had been gained pointing to the lack of competitiveness and some specific problems associated with this maize. As researchers and farmers became increasingly aware of the complex and inter-related problems, high hopes and optimism set forth earlier were greatly overshadowed affecting resource funding and a decline in research efforts. Many institutions either abandoned or drastically reduced their research involvement in such activities. As funding squeeze continued, only a few research centers and institutions could sustain needed research efforts in this area. The institutions which continued vigorously and persistently in improving the protein quality were International Center for Maize and Wheat Improvement (CIMMYT), the University of Natal, South Africa, and the Crow's Hybrid Seed Company in Milford, Illinois. All three centers had good research programs and were indeed quite successful in developing products with good agronomic performance and market acceptability.

Choosing among various options and strategies

As problems plaguing original soft opaque-2 materials became obvious and with no definite solution or particular strategy in sight at that time, many breeders and researchers really found themselves in a state of dilemma and confusion. This historical tragedy brought a turning point in the breeding efforts to take a new hard look examining

carefully the nature and seriousness of inherent problems. Equally important was to think of some pertinent options and strategies that could be deployed effectively in developing germplasm competitive in agronomic performance and free of problems considered important in acceptance. A detailed discussion of these options will not be attempted but rather salient features of interest will be summarized. Most options fall into two broad categories either involving or not involving specific mutant genes. The later group includes such options as recurrent selection for improved protein quality, altering germ-endosperm ratio, and increasing aleurone layers of the maize grain. All these approaches have been tried and require long-term efforts to achieve lysine levels matching the specific mutants which give a big boost in lysine content. In the former group are two interesting approaches. One of the approach exploits double mutant combination(s) whereas second one is based on the use of two genetic systems involving opaque-2 gene and the genetic modifiers of the opaque-2 locus.

The double mutant combination involving interaction of o2 and su2 has some advantages such as vitreous kernels, acceptable kernel appearance, lesser ear rot, increased lysine levels and better digestibility of protein. Yield is its only disadvantage which is sum total of independent negative effects of the two genes. Double mutant combination o2fl2 also has been researched and not pursued further as modified vitreous kernels are encountered in only a few genetic backgrounds. The most successful and rewarding option involves combined use of opaque -2 and the genetic modifiers of the o2 locus. Encouraging results from this strategy have been achieved at CIMMYT and elsewhere. The following sections will provide in-depth discussion on various aspects of this strategy which has been used extensively at CIMMYT.

Sequence of events and strategies in developing Quality Protein Maize

Groundwork and experimental evidence had to be accumulated on different breeding options before making an-allout shift in adopting a strategy which in the opinion and judgement of CIMMYT researchers will yield needed results. Several factors have contributed to the success of QPM work at CIMMYT but the single most important aspect in my opinion had been research strategy, methodology and constant adjustments from time to time based on needs. Laboratory support was a must in deploying this strategy to provide for rapid, and timely service in performing analyses. The interdisciplinary approach was another aspect of QPM research work which is considered exemplary. A program of this nature had to evolve continuously and as a consequence periodic adjustments and revision in program's philosophy, objectives and research methodology had to be made. Another aspect of this strategy which I consider imperative was tackling of complex and inter-related problems in a single multi-pronged strategy which in the end will result in a product that will have acceptance.

The approach CIMMYT researchers practiced involves combined use and interaction of two genetic systems; the simple opaque -2 system and a complex genetic modifier system. This approach has proved exceedingly successful and is being used in many institutions actively engaged in QPM breeding. This approach also has the advantage in that it attempts to solve key problems in an integrated manner. At this point a brief mention be made of research philosophy and some guiding principles in the use of this strategy. This approach emphasizes development of competitive QPM genotypes with quality protein as a bonus. To achieve this goal a conservative approach was used

initially with respect to biochemical characteristics. Since o2 gene boosts lysine levels to a two-fold increase, efforts were devoted to maintenance rather than further enhancing the levels of lysine at protein levels of 9-10% in the whole grain. This did facilitate greatly in breeding agronomically superior QPM genotypes focusing on critical and key problems.

Next I have attempted to describe sequence of everchanging and evolving QPM research. A bulk of high quality protein work in the initial stages involved conversion program as well as formation and improvement of soft endosperm opaque-2 populations and composites. As this work continued, meagre resources were allocated in doing exploratory work on finding appropriateness of alternative options and strategies. During the conversion programs, partially modified kernels had been observed by researchers at CIMMYT and elsewhere. Initially significance of such kernels, was not completely understood. In most instances the breeders tended to discard them considering as doubtful kernels. The first published report highlighting the importance of such kernels appeared in Crop Science journal in 1969 (Paez et al., 1969). Later such modified kernels have been observed by many researchers including CIMMYT breeders. Separation of such kernels when encountered began at CIMMYT as early as in 1969 by Dr. John Lonnquist and Dr. V.L. Asnani. Modified kernels were classified in different categories and sent to laboratory for analyses to study the effect of degree of modification on biochemical characteristics. A modest effort, primarily as an exploratory research, continued for a few years. Modified ears were sorted from the every possible source as they appeared during the conversion programs and during the process of seed increase of opaque-2 maize materials. As expected the modification was exceedingly poor ranging from slight to less than 25%. Also it may be important to mention that such kernels were encountered in different frequencies from different genetic backgrounds. Carribean, Cuban, and flint backgrounds in general tended to exhibit a higher frequency of modified kernels. Then as luck would have it, we identified a few opaque-2 converted populations and population crosses which had unusually higher frequency of modified kernels. Our hopes and enthusiasm from this point onwards took a different turn. A definite tilt and increasing resources were allocated in developing QPM germplasm. A series of actions followed this development.

Development of QPM donor stocks

The development of QPM donor stocks with well modified kernel phenotype and good protein quality was indeed quite important in accelerating rapid development of QPM germplasm. Initial effort was not only difficult but also frustrating. Selection for kernel modification had to be practiced at all stages while simultaneously maintaining protein quality. Intrapopulation selection was enforced for the accumulation of modifiers. In addition selection for improved kernel phenotype continued independently in selected modified opaque-2 families. Using these materials several promising materials from CIMMYT and from the national maize programs were converted to QPM. QPM donor stocks thus developed set a stage for the development of QPM germplasm in a wide array of genetic backgrounds. Basically two approaches were used in developing QPM donor stocks. The first approach was intrapopulation selection for genetic modifiers in opaque-2 backgrounds exhibiting a higher frequency of modified o2 kernels. Four tropical populations and one highland population which met this criteria were chosen for this

purpose. These populations were Composite K (H.E.o2), Ver.181-Ant.gpo2* Venezuela-1 o2, Thai o2 Composite, PD (MS6) H.E.o2, and Composite I. The breeding procedure used was controlled full-sib pollinations in the initial cycle followed by modified ear-to-row system suggested by Lonnquist (1964). Selection was practiced for modified ears and modified kernels at all stages.

The second approach involved recombination of superior hard endosperm opaque-2 families. The yellow and white families were recombined separately to develop Yellow H.E.o2 composite and White H.E.o2 composite respectively. Genetic mixing coupled with selection of modified ears and modified kernels of good protein quality was practiced for 3-4 cycles. By the mid-70s, it was felt that these materials had achieved a high frequency of modifiers and could be used as QPM donor stocks as well as QPM populations for further improvement using appropriate schemes.

Expanded efforts in QPM germplasm development

The development of QPM donor stocks led to a large scale QPM germplasm development efforts in different genetic backgrounds representing tropical, subtropical, and highland maize germplasm involving different maturities as well as grain color and texture. This was the beginning of an expanded effort in QPM germplasm development. Potentially useful normal maize populations were identified for QPM conversion program. Owing to the complexity and nature of modified phenotype trait, it was realized in the beginning that a standard backcross program will not work. An innovative breeding procedure was designed to handle the conversion process with the aim of obtaining QPM counterparts of normal genotypes as rapidly as possible. The procedure was named as a modified backcrossing -cum-recurrent selection program. The details of this procedure can be found in CIMMYT publications (Vasal, et al. 1980, 1984). A number of advanced maize populations in CIMMYT maize program were converted to QPM using this procedure. During conversion process emphasis was placed on yield, kernel modification and appearance, reduced ear rot incidence, rapid drying and agronomic attributes. Compared to normal counterparts, some of the QPM versions have given competitive performance in yield, and other agronomic traits (Tables 1 & 2). In addition to the conversion program described above, considerable resource allocation and research efforts were devoted to the formation and development of broadbased QPM gene pools. Several different ways were used in forming and improving these pools.

1. By genetic mixing of hard endosperm opaque-2 varieties, variety crosses, and also hybrids, if available.
2. Genetic mixing and recombination of soft opaque-2 materials followed by selection of genetic modifiers in homozygous o2 genetic backgrounds.
3. Crossing soft opaques with one or more hard endosperm opaque-2 populations with subsequent emphasis on selection for genetic modifiers in each cycle of selection.
4. Recombination of hard endosperm opaque-2 inbred or non-inbred families selected independently in each of the several o2 populations. Controlled pollination by hand or alternatively half-sib isolation can be used to achieve this objective.

5. Crossing of normal genotypes to one or more QPM donors followed by advancing F1 crosses to F2 by controlled sib-mating or in an isolation by planting balanced bulk and/or individual ears in a half-sib recombination block.

Amongst the procedures listed above, options 4th and 5th were mostly used in CIMMYT's work on QPM.

A total of seven tropical and six subtropical QPM gene pools were developed and improved continuously using a modified half-sib improvement selection system. The progress achieved in these pools is shown in figure 1. It may be pointed out that since most of the QPM gene pools were handled using non-inbred families, one may expect a low frequency of good lines surviving the inbreeding process.

In a period extending over 5-6 years, a huge volume of QPM germplasm was developed that could meet needs of several production environments in the tropical and subtropical areas. At this point it was realized that continuing conversion program will serve no useful purpose. Rather perhaps it will be more advantageous to accelerate progress by switching over to working in homozygous opaque-2 genetic backgrounds. The QPM germplasm consolidation and reorganization was considered appropriate at this stage.

New thrusts in QPM germplasm management strategy

As soon as QPM program succeeded in developing a huge volume of germplasm, it was important to think of the next strategy that will help in reducing germplasm volume and at the same time permit systematic and efficient handling of this valuable germplasm. Merging, consolidation, and reorganization of QPM germplasm was thus attempted. Knowledge of the germplasm as well color and maturity considerations were the principal guiding factors in this merging process. From the available QPM germplasm, eight tropical and six subtropical QPM gene pools were formed. These pools are listed in table 3.

At the same time ten QPM advanced populations were identified. Six of the populations were tropical and four subtropical in adaptation. These populations are listed in table 4 and were considered for international progeny testing trials (IPTT) for further improvement and as mechanism of disseminating germplasm to the NARs. Handling of QPM materials in homozygous opaque-2 background was emphasized at this point to facilitate faster progress and rapid accumulation of favorable modifiers for kernel modification, weight and density. Also working in homozygous o2 backgrounds had the additional advantage of reducing errors which generally occur due to misclassification and selection of kernels during the segregating generations. Selection in homozygous genetic backgrounds also permit choosing completely modified or normal looking kernels with certainty. The handling of QPM germplasm, both pools and populations, involving different stages have been discussed in several earlier publications from CIMMYT researchers (Vasal, et al. 1979,1978,1984,1997).

Progress in QPM germplasm

This section will describe briefly progress that has been attained in QPM germplasm for key traits that have been the stumbling block in gaining acceptance of these cultivars in the commercial production fields. It is encouraging to report that substantial progress has occurred in several materials. The yield gap has been narrowed down as can be seen in figure 2. Average kernel modification scores have improved with concomittant increase in the frequency of kernels with better modification scores (Figure 3).

Ear rot incidence in QPM materials has reduced gradually. Several materials are perhaps no different from the normal but still some show slightly higher incidence(Figure 4). In several studies reported earlier, no differences were encountered in normal and QPM counterparts for moisture content. In preliminary studies carried out in CIMMYT entomology laboratory, the QPM were no different from normal in damage by stored grained insect-pests. During all stages of improvement, the protein quality was monitored and maintained (Tables 5 & 6).

Emphasis on QPM hybrid development

The QPM hybrid initiative at CIMMYT was introduced in 1985. It offered obvious advantages including seed purity, monitoring and maintaining protein quality, enhanced modification and added uniformity and stability, enhanced yield potential, circumventing problems more efficiently and effectively, greater market acceptability, and more importantly helping growth of seed industry. Combining ability studies in QPM germplasm have been conducted and published (Vasal et al. 1993a, 1993b). Inbred line development efforts have been strengthened and evaluated for combining ability. Several hybrid combinations have been tested internationally and some of them have performed equal or better than some of the local checks included in the trials. (Tables 7 & 8). The performance of QPM hybrids is quite encouraging for adoption in some interested countries. Unfortunately due to funding constraints CIMMYT management had to drop completely QPM activities beginning 1993. The program has been re-initiated with full vigor and renewed enthusiasm. In the past 3-4 years, many more hybrid combinations have been tested again and the outlook of QPM appears quite promising. Because of this many countries in the developing world have become increasingly interested in QPM efforts.

Renewed emphasis on QPM in some national programs

In recent years several countries have shown unusual optimism and excitement in promoting and disseminating QPM cultivars to the farmers. Mass utilization of QPM variety-OBTANPA in Ghana is exemplary. More than 50% of the area in the country is planted to this variety. Several potential commercial channels for QPM utilization in Ghana have been identified including infant and institutional feeding programs and animal producers especially poultry and piggery farms. The Brazilian program has sustained QPM efforts all these years and have commercialized two QPM varieties BR451, and BR473. A double cross QPM hybrid has also been released in 1997. The program has greatly strengthened hybrid efforts and may soon be releasing more hybrids. The potential of QPM hybrids in China is tremendous. Also, QPM efforts are going on in

several academies all across the country. The Chinese government has shown great commitment to promote QPM hybrids all over the country. Only last month a QPM hybrid Zhongdan 9407 was released in a special ceremony attended by Vice Minister of Agriculture, President and Vice President of CAAS, DG CIMMYT, Mr. Sasakawa and his team from Japan, Dr. Borlaugh and several other CIMMYT and CAAS researchers. Field days and demonstrations have been held last year and this year. There is a new wave of enthusiasm to promote and disseminate hybrids. In Guizhou province (southern China) QPM intervention is being used to alleviate poverty and to improve economic well-being of the farmers. QPM interest in Mexico has grown with the support and commitment of Mexican government to cover substantial area under QPM hybrids in the next 2-3 years. A few countries in central America are ready to release QPM hybrids and to grow several thousands of hectares with such materials. India has also released Shakti-1 last year and is deeply involved in testing hybrids from CIMMYT. Several countries of Asia, Africa, and Latin America are part of QPM network on the improvement and promotion of Quality Protein Maize in selected developing countries.

In addition to the countries mentioned above, a number of other countries are also participating in the QPM network such as Honduras, Bolivia, Colombia, Ethiopia, Mozambique, Tanzania, Uganda, Zimbabwe, and south Africa. Several of these countries will be testing QPM varieties and hybrids and will also be involved in on farm trials, strip tests, and in conducting field days. The Republic of South Africa had earlier released hybrids HL-1, HL-2, and has recently released HL8 which has hard endosperm, good yield potential, and tolerance to diseases. The Sasakawa Global 2000 Project has helped to promote QPM in Ghana and is assisting some other countries in Africa in the promotion of QPM.

Conclusions and future outlook

The strategy used by CIMMYT researchers in developing QPM has proven to be successful and appears to be the only viable option that merits future consideration and use in research. Practically all QPM research programs in different countries are now using this approach based on the combined use of the opaque-2 gene and the genetic modifiers. Hybrid development efforts in QPM have become increasingly important as is evident from trials distributed in the past two years. It is hoped that many countries now involved in network will be able to select one or two most promising hybrids for release in their respective countries. More hybrid releases may be expected in China where hopes and optimism for QPM promotion and dissemination are exceedingly high. We may also witness increasing use of molecular genetic tools in QPM research in the future.

Programs at CIMMYT, Texas A&M, Brazil, and South Africa are already using these new tools on a limited scale. The mechanism controlling kernel modification needs to be further elucidated and the role of gamma zein to become more clear. Better characterization of zein proteins can perhaps facilitate and expedite conversion programs. In future perhaps more sophisticated techniques will become available that could distinguish modified $\alpha 2$ kernels from the normals.

References

- Bjarnason, M. and Vasal, S.K., Breeding of quality protein maize (QPM), in *Plant Breeding Rev.*, Janick, J., Ed., 1992, 181.
- Harpstead, D.D., High lysine maize in its proper perspective, in *Proc. 24th Annual Corn and Sorghum Research Conference*, American Seed Trade Assoc., Chicago, IL, 1969, 74.
- Ito.S., 1998. General Overview on Exploding Maize Demand in Asia. International Maize Conference in Tottori, June 1 to 4, 1998.
- Lonnquist, J.H. 1964. Modification of the ear-to-row procedure for the improvement of maize populations. *Crop Sci.* 4:227-228.
- Mertz, E.T., Bates, L.S., and Nelson, O.E., Mutant gene that changes protein composition and increases lysine content of maize endosperm, *Science*, 145, 279, 1964.
- Nelson, O.E., Mertz, E.T., and Bates, L.S., Second mutant gene affecting the amino acid pattern of maize endosperm proteins, *Science*, 150, 1469, 1965.
- Osborne, T.B. and Mendel, L.B., Amino acids in nutrition and growth, *J. Biol. Chem.*, 17, 325, 1914.
- Paez, A.V., Helm, J.L., and Zuber, M.S., Lysine content of opaque-2 maize kernels having different phenotypes, *Crop Sci.*, 9, 251, 1969.
- Vasal, S.K., Villegas, E., Bauer, R., Present status of breeding quality protein maize, in *Seed Protein Improvement in Cereals and Grain Legumes*, IAEA, Vienna, 1979, 127.
- Vasal, S.K., Villegas, E., Bjarnason, M., Gelaw, B., and Goertz, P., Genetic modifiers and breeding strategies in developing hard endosperm opaque-2 materials, in *Improvement of Quality Traits of Maize for Grain and Silage Use*, Pollmer, W.G. and Phipps, R.H., Eds., Nijhoff, The Hague, 1980, 37.
- Vasal, S.K., Villegas, E., and Tang, C.Y., Recent advances in the development of quality protein maize at the Centro Internacional de Mejoramiento de Maiz y Trigo, in *Cereal Grain Protein Improvement*, IAEA, Vienna, 1984^a, 167.
- Vasal, S.K., Villegas, E., Tang, C.Y., Werder, J., and Read, M., Combined use of two genetic systems in the development and improvement of quality protein maize, *Kulturpflanze*, 32, 171, 1984^b.
- Vasal, S.K. 1994. High quality protein corn. P. 80-121. In: A.R. Hallauer (ed.), *Speciality corns*. CRC Press, Boca Raton, FL.

Table 1: Across Location, Performance of Normal Materials and Corresponding Tropical QPM Germplasm, 1987

Material	Grain yield		QPM as percentage Of normal
	Normal	QPM	
	Kg ha-1		
Pool 23	5405	5330	98.6
Pool 24	5706	5457	95.6
Tropical High-oil	5733	5170	90.2
Population 62	5347	5484	102.6
Population 65	5255	5369	102.2
Population 63	5705	6236	109.3

Source: Bjarnason and Short, 1988.

Table 2: Mean Grain Yield of Four Highest Yielding QPM Varieties, Normal Reference Variety, and Best Local Checks At 11 Locations, CIMMYT EVT 15B, 1988

Variety	Grain yield Mt ha	Grain yield % of normal ref. entry
Across 8563 QPM	5.68	103.2
Capinopolis 8563 QPM	5.48	99.7
Iboperenda 8563 QPM	5.47	99.4
Tarapoto 8563 QPM	5.37	99.7
Suwan 8222 NRE	5.50	100.0
Best local check	5.36	97.5

Source: CIMMYT.

Table 3: QPM Gene Pools and Their Characteristics

QPM Pool Number	Adaptation	Maturity	Seed Color	Seed Texture
Pool 15 QPM	Tropical	Early	White	Flint-Dent
Pool 17 QPM	Tropical	Early	Yellow	Flint
Pool 18 QPM	Tropical	Early	Yellow	Dent
Pool 23 QPM	Tropical	Late	White	Flint
Pool 24 QPM	Tropical	Late	White	Dent
Pool 25 QPM	Tropical	Late	Yellow	Flint
Pool 26 QPM	Tropical	Late	Yellow	Dent
Pool 27 QPM	Subtropical	Early	White	Flint-Dent
Pool 29 QPM	Subtropical	Early	Yellow	Flint-Dent
Pool 31 QPM	Subtropical	Intermediate	White	Flint
Pool 32 QPM	Subtropical	Intermediate	White	Dent
Pool 33 QPM	Subtropical	Intermediate	Yellow	Flint
Pool 34 QPM	Subtropical	Intermediate	Yellow	Dent

Table 4: QPM Populations and Their Characteristics

Population number	Name	Adaptation	Maturity	Seed color	Seed texture
61	Early Yellow Flint QPM	Tropical	Early	Yellow	Flint
62	White Flint QPM	Tropical	Late	White	Flint-Semiflint
63	Blanco Dentado-1 QPM	Tropical	Late	White	Dent
64	Blanco Dentado-2 QPM	Tropical	Late	White	Dent
65	Yellow Flint QPM	Tropical	Late	Yellow	Flint
66	Yellow Dent QPM	Tropical	Late	Yellow	Dent
67	Templado Blanco Cristalino QPM	Subtrop.	Interm.	White	Flint
68	Templado Blanco Dentado QPM	Subtrop.	Interm.	White	Dent
69	Templado Amarillo QPM	Subtrop.	Interm.	Yellow	Flint
70	Templado Amarillo Dent QPM	Subtrop.	Interm.	Yellow	Dent

Table 5: Percent Protein, Tryptophan, and Lysine in the Whole Grain of QPM Pools

QPM Gene Pool	Protein %	Tryptophan in protein %	Lysine in protein %	Quality index
Pool 15 QPM	9.1	0.94	4.2	4.6
Pool 17 QPM	8.9	1.04	4.5	4.5
Pool 18 QPM	9.9	0.93	4.0	4.6
Pool 23 QPM	9.1	1.03	3.8	4.2
Pool 24 QPM	9.4	0.92	3.8	4.0
Pool 25 QPM	9.8	0.94	4.0	4.0
Pool 26 QPM	9.5	0.90	4.1	4.3
Pool 27 QPM	9.5	1.05	4.2	4.8
Pool 29 QPM	9.2	1.06	4.3	4.8
Pool 31 QPM	10.2	0.96	4.1	4.5
Pool 32 QPM	8.9	1.04	4.2	4.5
Pool 33 QPM	9.3	1.05	-	4.2
Pool 34 QPM	9.1	1.10	4.1	4.5

Table 6: Protein Quality in Different Cycles of Selction
(Endosperm analysis)

Material	Cycle	Protein (%)	Tryptophan (%)	Lysine (%)
CIMMYT H.E. o2	C0	8.0	0.78	2.90
	C4	8.6	0.77	2.90
PD (MS) 6 H.E. o2	C0	8.5	0.78	2.50
	C4	8.8	0.78	2.60

Table 7: Superior white QPM hybrids tested across fifteen locations at El Salvador, Guatemala and Mexico, 1998.

Pedigree	Yield t/ha	Ear rot (%)	Tryptophan (%)	Ear Modification	Silking (Days)	Plant height (cm)
CML142xCML146	6.48	3.7	0.096	2.0	55	242
CML159xCML144	6.39	4.3	0.100	1.6	56	230
(CL Q6203xCML150)CML176	6.28	5.7	0.088	2.1	55	239
CML145xCML144	5.81	5.8	0.840	2.0	54	241
CML158xCML144	5.59	7.1	0.103	1.3	55	228
CML146xCML150	5.48	8.7	0.084	3.6	56	222
POZA RICA 8763 TLWD	5.34	12.0	0.095	2.8	54	230
Normal hybrid check	5.58	9.5	0.070	2.0	56	228
Local checks: HB-83, CB-HS-5G, H-59, XM7712, GUAYOPE						

Table 8: Superior subtropical yellow QPM hybrids tested across five locations at Mexico and Zimbabwe, 1998.

Pedigree	Yield t/ha	Ear rot (%)	Ear Modification	Silking (Days)	Plant height (cm)
CML161xCML165	9.82	0.1	1.5	98	219
CLQ3301xCML193	8.09	6.0	2.0	91	207
CLQ6901xCML193	8.08	2.1	1.5	94	216
CLQ3302xCML193	7.78	3.3	1.5	92	198
CLQ3401xCML193	7.78	1.5	2.0	93	199
CLQ3402xCML193	7.69	2.0	2.5	94	197
Normal hybrid check	9.82	1.8	4.5	98	223
Local checks: H-358, A-7595, H-312					